



Capsular fixation limits graft extrusion in lateral meniscal allograft transplantation

Angel Masferrer-Pino¹ · Joan C. Monllau^{1,2} · Ferran Abat³ · Pablo E. Gelber^{1,4}

Received: 3 June 2019 / Accepted: 15 August 2019
© SICOT aisbl 2019

Abstract

Purpose The main purpose of this investigation was to compare the amount of graft extrusion of lateral meniscal allograft transplantation (MAT) performed with a suture-only technique with or without a capsulodesis. Secondly, the assessment of functional results was also covered. We hypothesized that capsular fixation reduces the post-operative degree of allograft extrusion and it does not affect the functional outcomes during the short-term follow-up period studied.

Methods Prospective series of 29 lateral MAT. Fifteen were fixed with a suture-only technique (group A). The remaining 14 cases (group B) also included arthroscopic lateral capsular fixation (capsulodesis). Functional results were assessed with Lysholm, Tegner, and VAS for pain. Magnetic resonance imaging (MRI) was performed to determine the degree of meniscal extrusion. Millimeters of extrusion and percentage of extruded meniscal tissue were calculated for both groups. The degree of extrusion was considered minor if it was < 3 mm or major if it was > 3 mm.

Results Group A had 11 cases (73.3%) of major extrusion and group B had 4 cases (28.6%) ($p = 0.02$). The percentage of extruded meniscal tissue was 35% in group A and 24.6% in group B ($p = 0.04$). At a mean 3.4 years (range 1–4) post-operatively, the Lysholm score had a mean 89.60 ± 6.93 and 91.43 ± 6.19 points in groups A and B, respectively ($p < 0.001$). The median follow-up Tegner score improved from 4 (range 3–5) to 7 (range 6–9) in group A ($p < 0.001$) and from 4 (range 3–5) to 7 (range 6–8) in group B ($p < 0.001$). VAS dropped 5 and 7.3 points in groups A and B, respectively ($p < 0.001$). There were no complications in this series.

Conclusions In lateral MAT with the suture-only fixation technique, the described capsulodesis minimized meniscal extrusion. In terms of functional results, there were no differences between the groups at a mean 3.4-year follow-up.

Keywords Meniscal transplantation · Meniscal extrusion · Meniscus · Knee · Capsulodesis

Introduction

Meniscal allograft transplantation (MAT) was developed in Germany in the mid-1980s [1]. This procedure replaces the lost meniscal tissue to eventually prevent progressive deterioration

of the joint due to a previous meniscectomy [2, 3]. To address the problems of meniscectomized patients, there are ongoing efforts to develop techniques for meniscus regeneration or meniscus scaffolding using tissue engineering strategies [4, 5]. Progressively, MAT has become a valid treatment option in relatively young and active patients symptomatic due to a prior meniscectomy [6]. It is currently accepted that MAT provides favorable clinical results on at least the short- and medium-term basis [7]. Several soft tissue and bone fixation surgical techniques have been described to better fix the graft and to prevent mechanical failures. However, no significant clinical differences have been reported between them to date [7]. Extrusion is an interesting phenomenon that is usually identified shortly after transplantation and seems to be stable over time [8]. It is defined as a tendency toward radial displacement of the graft beyond the tibial margin and has been often observed after MAT [9]. Biomechanically, an extruded meniscus would

✉ Pablo E. Gelber
personal@drgelber.com

¹ Department of Orthopedic Surgery and Traumatology, ICATME, Hospital Universitari Dexeus - Universitat Autònoma de Barcelona, Sabino Arana, 5-19, 08028 Barcelona, Spain

² Department of Orthopaedic Surgery, Parc de Salut Mar – Universitat Autònoma de Barcelona, Barcelona, Spain

³ ReSport Clinic, Barcelona, Spain

⁴ Department of Orthopaedic Surgery, Hospital de la Santa Creu i Sant Pau - Universitat Autònoma de Barcelona, Barcelona, Spain

decrease the resistance to hoop strains. The MAT with suture-only fixation has shown a greater degree of extrusion when compared to bony fixation [10, 11]. Although extrusion has not shown to have any clinical consequences yet, the abnormal position of those grafts causes concern among surgeons. Therefore, different strategies have been suggested to limit or prevent MAT extrusion. Recently, a technique that aims to stabilize the residual meniscus rim or the lateral capsule to the tibial plateau (capsulodesis) has been implemented to reduce extrusion in lateral MAT [12].

The main purpose of this investigation was to compare the amount of graft extrusion of lateral MAT performed with a suture-only technique with or without a capsulodesis. Secondly, the assessment of functional results was also covered. We hypothesized that capsular fixation reduces the post-operative degree of allograft extrusion and it does not affect the functional outcomes during the short-term follow-up period studied.

Methods

A randomized and prospective clinical trial was performed following the CONSORT guidelines. The study protocol was approved by the institutional review board (*ExtMen 2016-01*). Informed consent was signed from each patient. All the surgical procedures were performed by two of the authors. Twenty-nine consecutive patients were operated on with a lateral MAT. The fixation technique in each patient was randomly determined with permuted blocks for the suture-only group (A) or the capsulodesis group (B).

In all the cases, MAT was indicated for lateral joint line pain due to a previous large meniscectomy that had not improved with non-surgical therapies. Although no relationship between malalignment and meniscal extrusion has been established, patients with alignment beyond 5 degrees of varus or valgus were excluded. Patients who had an Ahlback grade greater than II were also excluded. Neither were cases performed with any concomitant surgical procedure or patients with a body mass index greater than 30 considered for this study.

Radiographic and functional evaluations were performed by two independent observers (both, orthopaedic surgeons). With regard to the functional evaluation, the observers were blinded to the type of MAT fixation used.

Surgical technique

Allograft sizing was matched with the donor's morphometric dimensions (weight and size) and double-checked with the method described by Pollard et al. [13]. The allografts were fresh-frozen, non-irradiated and non-antigen matched. They were provided by an authorized local tissue bank.

The whole surgical technique was performed arthroscopically in all the cases. The specific details of the suture-only [10, 14] as well as the capsulodesis techniques have been previously described [12, 15]. In all the procedures, the remains of the host menisci were assessed and refreshed using a combination of shaving, rasping, and high-frequency trephination to promote healing [16]. During the surgical procedure, all the allografts were soaked and kept in a vancomycin-saline solution at 36 °C [17]. In all the cases, both allograft horns were fixed to 4.5-mm transtibial tunnels drilled at the anatomic anterior and posterior meniscal root attachments [18]. Once the allograft was correctly placed, final fixation was accomplished with a combination of all-inside (FasT-Fix; Smith & Nephew, Andover, MA) vertical or horizontal sutures in the posterior half of the meniscus, up to the popliteus hiatus, and with outside-in sutures in the anterior half. The sutures were placed every 5 to 10 mm alternating on the superior and inferior aspects of the graft to obtain a better end-position of the MAT and improve its fixation to the original rim or capsule. At the end of the procedure, the sutures placed on the horns were tied together over the tibial cortex.

All the allografts that were fixed with only transosseous sutures in the anterior and posterior horns were considered as group A. In group B, prior to the graft transplantation, the lateral meniscal rim or capsule was also fixed to the lateral tibial plateau margin. The redundant or loose lateral capsule was first identified. If any osteophyte was observed on the lateral edge of the tibial plateau, a motorized burr was used to remove it. Two 2.4-mm tunnels were drilled from the antero-medial tibial cortex aiming to the edge of the lateral tibial plateau, leaving a bridge of 10 mm between them, where the capsule was seen more laterally displaced (Fig. 1). It was done with the help of a tibial ACL guide (Pinn-ACL Guide, ConMed, Largo, Florida).

One suture was then passed through each tunnel. The capsule including any meniscal remnant was pierced with an 18-gauge spinal needle loaded with a #2 PDS suture using an outside-in technique. These shuttle sutures were replaced by high strength sutures and pulled down through each tibial tunnel. This maneuver brings the capsule over the LTP, reducing any redundancy (capsulodesis). Finally, the two strands were tied to each other on the tibial cortex, similarly to the meniscal horn fixation (Fig. 2). Alternatively, they can be tied over a button. Once the capsulodesis had been done, the meniscal transplantation itself was performed.

Post-operative care

All the patients had the same post-operative protocol. The first two weeks, only proprioceptive weight-bearing with a knee brace blocked in full-extension was allowed. Progressive weight-bearing was subsequently encouraged. Full weight-

bearing was regularly obtained by week four to six, depending on the patients' tolerance. Range of motion exercises was limited to 90 degrees the first four weeks, increased progressively by week six and then to full range of motion afterwards. At four months of index surgery, most of the patients returned to normal activity and workloads.

Magnetic resonance imaging evaluation

An MRI was performed on each patient in the supine position before surgery and between the 12 and 36-month follow-up. All the MRIs were done in a 1.5-T superconducting magnet (Prestige 2T, Elscint, Haifa, Israel) using a knee-specific circular coil. The protocol for each study was axial fast spin echo T2-weighted with fat saturation, coronal fast spin echo intermediate-weighted, sagittal spin echo intermediate-weighted, and sagittal fast spin echo T2-weighted with fat saturation. The images were evaluated twice, three weeks apart. Two independent observers (orthopaedic surgeons) assessed the images. The mean of these two measurements was used in the analysis. The PACS workstation (Centricity Enterprise Web V3.0, General Electric Healthcare, Milwaukee, Wisconsin) was used for the study. As reported, the graft position was evaluated on coronal images [16], where extrusion is maximum. In general, those images were at the level of the medial collateral ligament. Measurement was performed drawing two lines. One of them was a vertical line intersecting the peripheral margin of the LTP at the point of transition from horizontal to vertical. A perpendicular line was also drawn from the outer margin of the meniscus to the former line.




Fig. 1 Left knee, antero-medial view. Lateral capsulodesis tunnels placement using the ACL pin guide through the antero-lateral (AL) portal




Fig. 2 Capsular fixation suture passed through transtibial tunnel and tied to each other on the medial cortex fixing the capsule to the tibial plateau

The latter was specifically done to measure the degree of extrusion [19, 20]. Additionally, when the graft was extruded less than 3 mm beyond the LTP, it was considered minor extrusion [20]. Conversely, major extrusion was considered when the allograft exhibited more than 3 mm of subluxation [20]. The percentage of extruded meniscal tissue was also calculated. Both the percentage of extrusion as well as it being classified as either major or minor extrusion were then compared between groups.

Functional evaluation

Before surgery and at the final follow-up, clinical outcomes were evaluated with the Lysholm score as well as Tegner Activity Scale. A 10-point visual analogical scale (VAS) for pain was also used. Patient satisfaction with a subjective score was also used. It was graded as very satisfied (4 points), satisfied (3 points), neutral (2 points), somewhat dissatisfied (1 point), and not satisfied at all (0 points). All the results were used to compare outcomes between groups.

Data analysis

The sample size was based on a priori power calculations for extrusion. While the categorical variables are reflected in percentages and frequencies, the continuous variables are expressed by a mean \pm standard deviation. The interobserver agreement was included using the intraclass correlation coefficient [21]. The values were interpreted based on previous studies [22]. A 95% confidence interval was calculated for all possible variables. Student's *t* test was used for independent

data with a statistical power of 80% and an alpha error of 0.05 to calculate the number of patients needed in each group to detect statistically significant minimum differences of 10% in relation to the degree of meniscal extrusion. There was a standard deviation of 3 and a maximum follow-up loss of 10%. The results obtained using Fisher's exact test to compare two samples leads us to think that it is broad enough for the analysis of the hypotheses of this study. Student's *t* test was also used to compare meniscal extrusion rates and functional scores between the two groups. The skewness and kurtosis test demonstrate that all the variables followed normal distributions.

All data analysis was assessed using the SPSS 19 package (SPSS Inc., Chicago, Illinois). Statistical significance was set at .05.

Results

The average radiological follow-up period with postoperative MR was 1.5 years (range, 1–3), while it was 3.2 years (range, 1–4) for clinical follow-up. All the patients were available for assessment. The series had 19 men (65.5%) and ten women (34.5%) with an average age of 40.9 years (range, 26 to 54 years). Sixteen (55.2%) MAT were performed in right knees while 13 (44.8%) were performed in left knees. There were 15 cases in group A (51.7%) and 14 cases in group B (48.3%). Table 1 shows that both groups were comparable in terms of radiographic findings, gender, age, and the pre-operative functional status.

Graft extrusion

In accordance with the extrusion criteria described above, 14 knees (48.3%) showed graft extrusion as minor (< 3 mm) while major extrusion (> 3 mm) was observed in 15 patients (51.7%). Within group A, there were four cases (26.7%) of minor extrusion and 11 cases (73.3%) of major extrusion. In group B, there were major and minor extrusion in ten (71.4%) and four cases (28, 6%), respectively (Table 2) ($p = 0.02$).

If we focus on comparing the degree of meniscus extrusion with the type of fixation used, a lower percentage of extrusion was also observed in group B. In group A, this percentage was 35 ± 9.8 . Conversely, $24.6\% \pm 15.5$ of extruded meniscal tissue was seen in group B ($p = 0.04$) (Table 3).

Clinical outcomes

A significant improvement in all the assessed scores was observed when the pre- and post-operative values were compared, regardless of the technique used. The Lysholm score improved from 62.3 ± 11.8 and 48.8 ± 13.9 to 89.6 ± 6.9 ($p < 0.001$) and 91.4 ± 6.19 ($p < 0.001$), in groups A and B, respectively (Fig. 3a). Similarly, the mean VAS score

decreased from 7.3 ± 2.1 to 1.3 ± 1.40 in group A ($p < 0.001$) and from 8.2 ± 1 to 0.9 ± 1 in group B ($p < 0.001$) (Fig. 3b). The Tegner activity score also improved from 4 (range, 3–5) to 7 (range, 6–9) in group A ($p < 0.001$) and from 4 (range, 3–5) to 7 (range, 6–8) in group B ($p < 0.001$) (Fig. 3c).

Patient satisfaction with the surgery with a maximum of 4 was 3.4 ± 0.3 points in group A and 3.8 ± 0.2 in group B. No differences were observed when comparing the functional variables analyzed between the two study groups. The intraclass correlation coefficient was classified as excellent (0.89, 95% CI 0.81 to 0.95). Finally, no complications have been reported in this series.

Discussion

The main finding of this investigation was that capsulodesis decreased the extrusion of a lateral MAT fixed with a suture-only technique. This confirmed the first hypothesis. Secondly, in terms of function, both groups obtained similar and comparable results in the short term. That fact confirms the second hypothesis of the study.

Several investigations have shown that transplanted menisci tend to extrude more than native menisci [23, 24]. Despite this extrusion, most series have shown good clinical results. Many aspects of meniscal extrusion are still unknown. Regarding the cause, it could be multifactorial. However, it remains unanswered as to whether this abnormal position may cause any alteration of the meniscus' biomechanics or not. However, concern exists that the non-anatomical position of the MAT might likely produce further damage to the knee in the long-term. The extrusion phenomenon seems to occur early after transplantation (< 6 weeks) and does not worsen posteriorly [8]. Similarly, those that do not extrude early are unlikely to further extrude [8]. The vast majority of series that deal with this topic investigated the radial displacement of the graft only on the coronal plane. Few authors have assessed MAT extrusion in an anterior or posterior direction [25, 26], which may also be significant in terms of final meniscal behaviour.

Several different strategies have been proposed to limit or minimize extrusion. Although the final cause is unknown, it is quite common to see a very loose and redundant capsule in lateral symptomatic meniscectomized compartments. That is probably the reason that the capsulodesis technique has been so effective at minimizing extrusion in the current series when compared to the otherwise widely used only-suture technique [10].

As mentioned earlier on, the cause of meniscal extrusion could be multifactorial. The size of the graft [24], as well as the mismatching of recipient and donor [13, 18], the compartment location (medial versus lateral) [26], and the type of fixation [10, 12, 14], all have been suggested as possible

Table 1 Composition of both groups before surgery

Variable	Group A	Group B	Significance (P)
Age, years	41 ± 8.5 (33.8, 41.7)	40.9 ± 7.0 (36.8, 45)	0.74
Gender, male/female, %	60/40	71.5/28.5	0.55
Lysholm	62.3 ± 11.8 (56.3, 68.4)	48.79 ± 13.90 (40.7, 56.8)	0.07
Tegner	4 (3–5) (2.96, 4.64)	4 (3–5) (2.95, 4.5)	0.87
Visual analog scale	7.3 ± 2.1 (5.17, 7.8)	8.21 ± 1 (7.6, 8.8)	0.51
Rx joint space narrowing, mm	3.1 ± 1.4 (2.62, 3.6)	3 ± 1.2 (2.8, 4.0)	0.33

Values expressed as mean, standard deviation, and 95% CI unless otherwise indicated

factors related to graft extrusion. With regard to the graft fixation method, investigated in the current study, it may influence both the failure rate and the kinematics of the knee joint. It has been shown, biomechanically, that extrusion might cause abnormal load transmission and later chondral impairment [27, 28]. It is likely that the extruded menisci tend to fail due to the incongruence with the femoral condyle. Therefore, maintaining the correct position of the allograft seems crucial to restoring the normal kinematics of the operated knee. While fixation of the graft meniscal body to the capsule or the meniscal remnant is usually performed with a standard meniscal repair technique, meniscal horn fixation is much trickier. Horn attachment is a critical step that requires strong fixation. In lateral MAT, to that end, the two most commonly used techniques are transtibial fixation with either only soft tissue or with a bone block in the allograft meniscal horns or the bone bridge in a trough technique [29]. Some authors sustain that bony fixation technique is superior to the only-suture technique in terms of restoring the normal biomechanics of the compartment and having fewer complications [30, 31]. However, MAT without bone fixation has also shown good and excellent results in terms of pain relief and clinical and functional results [23, 32, 33], allowing for a return to sports activities in top-level athletes [34, 35]. As has been recently shown, the graft also exhibits less extrusion when compared to bony fixation if capsulodesis is added to the lateral MAT fixed without bone plugs [12].

MAT extrusion is a phenomenon that appears often after transplantation [10, 12, 14, 26, 29, 32, 37–39], and has no

relationship with the pre-operative position of the meniscal remnant [36]. It is thought that a strong bony fixation technique keeps the meniscal graft in place and so tends to decrease the rate of extrusion even though that better positioning has not been related to clinical, functional, or radiological differences when compared to soft tissue fixation [10, 16]. As far as we know, the current study is the first comparing two different methods of MAT fixations without bone plugs. This may also be significant because some tissue banks worldwide only deliver meniscal allografts without bone blocks.

To date, different methods have been described to reduce the degree of MAT extrusion. Jang et al., in his series [24], suggested reducing the size of the graft by 5%. That approach makes for a reduction in the percentage of meniscal extrusion without producing any deterioration at the clinical or radiological level. When using a bone-bridge technique, the greater the obliquity of the trough on the axial plane, the more likely the risk of graft extrusion [38]. Therefore, assuring a correct starting point of the bone trough diminishes the angle and so the tendency of the MAT to laterally dislocate. Moreover, a simple resection of an osteophyte greater than 2 mm in the lateral tibial plateau can also reduce the extrusion of lateral MAT [39]. The rehabilitation process may even influence the degree of graft extrusion [37]. When compared to standard rehabilitation, delayed rehabilitation showed less coronal graft extrusion and joint space narrowing on weight-bearing and reduced the progression of degenerative changes. Lastly, to limit graft subluxation, some surgeons have suggested the fixation of the meniscal graft to the tibial plateau. Although that technique can be effective, it has the inherent risk of reducing the normal mobility of the meniscus during knee motion, particularly in the lateral compartment where the meniscus presents greater extrusion. Therefore, that technique

Table 2 Frequency of graft extrusion

	Minor	Major
Suture-only	4 (26.7%)	11 (73.3%)
Capsulodesis	10 (71.4%)	4 (28.6%)
Total	14 (48.3%)	15 (51.7%)

$p = 0.02$

Table 3 Graft extrusion percentage

Group	Observations	Mean	S.D.
Suture-only	15	35	9.8
Capsulodesis	14	24.6	15.5
p value	0.04		




Fig. 3 Overall improvement of functional scores in group B: **a** Lysholm, **b** VAS, **c** Tegner

was refined to preserve the meniscal motion by fixing the peripheral capsule or meniscal rim to the tibial plateau instead. Therefore, the MAT is subsequently secured to a firm lateral

wall that has been freshened beforehand to promote healing and graft incorporation [40]. This technique emphasizes the role that the fixation of the capsule could have on the tibial plateau. However, those authors used metallic anchors to fix the graft and failed to provide nor radiological neither clinical outcomes [40].

Despite the efforts to reduce graft extrusion after lateral MAT, there are still pre- and postoperative factors related to graft extrusion. Although the native lateral meniscal roots have been well studied [41], some of their peripheral and meniscal body attachments are still relatively unknown. Some authors consider that the lateral meniscotibial ligament (MTL) plays an important role in stabilizing the body of the lateral meniscus [42]. It could then be hypothesized that extrusion of the graft is to some extent due to the lack of MTL reconstruction during the MAT procedure. Thus, the current capsulodesis technique, with the tunnels drilled at the level of the anatomical attachments of the MTL, is somehow reproducing its function and so limiting the extrusion of the allograft [12]. In terms of extrusion, the proposed capsulodesis added to a soft tissue fixation through bone tunnels positively compares to bony fixation methods [12].

The described technique may thus provide additional biomechanical benefits to the MAT procedure. It is a versatile technique that avoids additional hardware while preserving good bone stock (2.4-mm transosseous tunnels). It also




Fig. 4 MRI at 6 months post-operatively showing the position of the mid-body of lateral meniscus allograft and the lateral capsulodesis tibial tunnel placement. In the superior right quadrant, arthroscopic view of the transplanted allograft performed after the capsulodesis

provides a satisfactory option to reduce lateral MAT extrusion when the graft is supplied by the tissue bank without bone blocks. Last but not least, being an implant-free technique allows for optimal postoperative MRI assessment (Fig. 4).

Although there is still a big dilemma regarding the impact of the meniscal extrusion on the future of the knee joint, this series showed one of the lowest meniscal extrusion rates to date [7, 9, 10, 12, 14, 26, 29, 32, 37–39]. Longer follow-up studies are needed to confirm these results. Anatomical studies are also required to further study the anatomy and biomechanics of the knee lateral structures and their impact on knee kinematics after MAT.

Different limitations to this investigation need to be addressed. A larger sample size could have reflected some differences in the clinical outcomes and thus, the possibility of a type II or beta error cannot be excluded. Also, differences in the clinical results could have been different if a longer follow-up had been carried out. However, the main objective of the study was to assess the degree of meniscal extrusion. It is a phenomenon that occurs in the first weeks after the surgery [8]. The intra-observer agreement of the MRI measurement method was not calculated.

Conclusions

In the lateral MAT suture-only fixation technique, the described capsulodesis minimized meniscal extrusion. In terms of functional results, there were no differences between groups at a mean 3.4-year follow-up.

Acknowledgments The authors thank Julio Florentino Del Corral for assisting in the statistical analysis as well as Eric Goode for his help in correcting the manuscript.

References

- Milachowski KA, Weismeier K, Wirth CJ (1989) Homologous meniscus transplantation: experimental and clinical results. *Int Orthop* 13(1):1–11
- DeHaven KE (1999) Meniscus repair. *Am J Sports Med* 27(2):242–250
- Zhang P, Zhao Q, Shang X, Wang Y (2018) Effect of arthroscopic resection for discoid lateral meniscus on the axial alignment of the lower limb. *Int Orthop* 42(8):1897–1903
- Verdonk R, Madry H, Shabshin N, Dirisamer F, Peretti GM, Pujol N, Spalding T, Verdonk P, Seil R, Condello V, Di Matteo B, Zellner J, Angele P (2016) The role of meniscal tissue in joint protection in early osteoarthritis. *Knee Surg Sports Traumatol Arthrosc* 24(6):1763–1774
- Xu C, Zhao J (2015) A meta-analysis comparing meniscal repair with meniscectomy in the treatment of meniscal tears: the more meniscus, the better outcome? *Knee Surg Sports Traumatol Arthrosc* 23(1):164–170
- Kim J-H, Lee S, Ha DH, Lee SM, Jung K, Cho W (2018) The effects of graft shrinkage and extrusion on early clinical outcomes after meniscal allograft transplantation. *J Orthop Surg Res* 13(1):181
- Rosso F, Bisicchia S, Bonasia DE, Amendola A (2015) Meniscal allograft transplantation: a systematic review. *Am J Sports Med* 43(4):998–1007
- Lee DH, Kim TH, Lee SH, Kim CW, Kim JM, Bin S (2008) Evaluation of meniscus allograft transplantation with serial magnetic resonance imaging during the first postoperative year: focus on graft extrusion. *Arthroscopy* 24(10):1115–1121
- Samitier G, Alentorn-Geli E, Taylor DC, Rill B, Lock T, Moutzouros V, Kolowich P (2015) Meniscal allograft transplantation. Part 1: systematic review of graft biology, graft shrinkage, graft extrusion, graft sizing, and graft fixation. *Knee Surg Sports Traumatol Arthrosc* 23(1):310–322
- Abat F, Gelber PE, Erquicia JI, Pelfort X, Gonzalez-Lucena G, Monllau JC (2012) Suture-only fixation technique leads to a higher degree of extrusion than bony fixation in meniscal allograft transplantation. *Am J Sports Med* 40(7):1591–1596
- De Coninck T, Huysse W, Verdonk R, Verstraete K, Verdonk P (2013) Open versus arthroscopic meniscus allograft transplantation: magnetic resonance imaging study of meniscal radial displacement. *Arthroscopy* 29(3):514–521
- Masferrer-Pino A, Monllau JC, Ibáñez M, Erquicia JI, Pelfort X, Gelber PE (2018) Capsulodesis versus bone trough technique in lateral meniscal allograft transplantation: graft extrusion and functional results. *Arthroscopy* 34(6):1879–1888
- Pollard ME, Kang Q, Berg EE (1995) Radiographic sizing for meniscal transplantation. *Arthroscopy* 11(6):684–687
- Abat F, Gelber PE, Erquicia JI, Tey M, Gonzalez-Lucena G, Monllau JC (2013) Prospective comparative study between two different fixation techniques in meniscal allograft transplantation. *Knee Surg Sports Traumatol Arthrosc* 21(7):1516–1522
- Monllau JC, Ibáñez M, Masferrer-Pino A, Gelber PE, Erquicia JI, Pelfort X (2017) Lateral capsular fixation: an implant-free technique to prevent meniscal allograft. *Arthrosc Tech* 6(2):e269–e274
- Furumatsu T, Miyazawa S, Fujii M, Tanaka T, Kodama Y, Ozaki T (2019) Arthroscopic scoring system of meniscal healing following medial meniscus posterior root repair. *Int Orthop* 43(5):1239–1245
- Pérez-Prieto D, Torres-Claramunt R, Gelber PE, Shehata TM, Pelfort X, Monllau JC (2016) Autograft soaking in vancomycin reduces the risk of infection after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 24(9):2724–2728
- Perez-Blanca A, Prado Nóvoa M, Lombardo Torre M, Espejo-Reina A, Ezquerro Juanco F, Espejo-Baena A (2018) The role of suture cutout in the failure of meniscal root repair during the early post-operative period: a biomechanical study. *Int Orthop* 42(4):811–818
- Puig L, Monllau JC, Corrales M, Pelfort X, Melendo E, Caceres E (2006) Factors affecting meniscal extrusion: correlation with MRI, clinical, and arthroscopic findings. *Knee Surg Sports Traumatol Arthrosc* 14(4):394–398
- Costa CR, Morrison WB, Carrino JA (2004) Medial meniscus extrusion on knee MRI: is extent associated with severity of degeneration or type of tear? *Am J Roentgenol* 183(1):17–23
- Hale CA, Fleiss JL (1993) Interval estimation under two study designs for kappa with binary classifications. *Biometrics* 49(2):523–534
- Landis JR, Koch GG (1977) The measurement of observer agreement for categorical data. *Biometrics* 33(1):159–174
- Verdonk PC, Demurie A, Almqvist KF, Veys EM, Verbruggen G, Verdonk R (2005) Transplantation of viable meniscal allograft. Survivorship analysis and clinical outcome of one hundred cases. *J Bone Joint Surg Am* 87(4):715–724
- Jang SH, Kim JG, Ha JG, Shim JC (2011) Reducing the size of the meniscal allograft decreases the percentage of extrusion after meniscal allograft transplantation. *Arthroscopy* 27(7):914–922

26. Kim NK, Bin SI, Kim JM, Lee CR, Kim JH (2017) Meniscal extrusion does not progress during the midterm follow-up period after lateral meniscal transplantation. *Am J Sports Med* 45(4):900–908
27. Lee DH, Lee CR, Jeon JH, Kim KA, Bin SI (2015) Graft extrusion in both the coronal and sagittal planes is greater after medial compared with lateral meniscus allograft transplantation but is unrelated to early clinical outcomes. *Am J Sports Med* 43(1):213–219
28. Chen MI, Branch TP, Hutton WC (1996) Is it important to secure the horns during lateral meniscal transplantation? A cadaveric study. *Arthroscopy* 12(2):174–118
29. Kim JG, Lee YS, Bae TS, Ha JK, Lee DH, Kim YJ, Ra HJ (2013) Tibiofemoral contact mechanics following posterior root of medial meniscus tear, repair, meniscectomy, and allograft transplantation. *Knee Surg Sports Traumatol Arthrosc* 21(9):2121–2125
30. Matava MJ (2007) Meniscal allograft transplantation: a systematic review. *Clin Orthop Relat Res* 455:142–157
31. Alhalki MM, Howell SM, Hull ML (1999) How three methods for fixing a medial meniscal autograft affect tibial contact mechanics. *Am J Sports Med* 27(3):320–328
32. Paletta GA, Manning T, Snell E, Parker R, Bergfeld J (1997) The effect of allograft meniscal replacement on intraarticular contact area and pressures in the human knee. *Am J Sports Med* 25(5):692–698
33. Alentorn-Geli E, Seijas R, García M, Alvarez P, Steinbacher G, Cuscó X, Vilarrubia R, Cugat R (2011) Arthroscopic meniscal allograft transplantation without bone plugs. *Knee Surg Sports Traumatol Arthrosc* 19(2):174–182
34. González-Lucena G, Gelber PE, Pelfort X, Tey M, Monllau JC (2010) Meniscal allograft transplantation without bone blocks: a 5- to 8-year follow-up of 33 patients. *Arthroscopy* 26(12):1633–1640
35. Alentorn-Geli E, Vazquez RS, Diaz PA, Cuscó X, Cugat R (2010) Arthroscopic meniscal transplants in soccer players: outcomes at 2- to 5-year follow-up. *Clin J Sport Med* 20(5):340–343
36. Chalmers PN, Karas V, Sherman SL, Cole BJ (2013) Return to high-level sport after meniscal allograft transplantation. *Arthroscopy* 29(3):539–544
37. Lee BS, Bin SI, Kim JM, Kim JH, Lim EJ (2017) Meniscal allograft subluxations are not associated with preoperative native meniscal subluxations. *Knee Surg Sports Traumatol Arthrosc* 25(1):200–206
38. Lee DW, Lee JH, Kim DH, Kim JG (2018) Delayed rehabilitation after lateral meniscal allograft transplantation can reduce graft extrusion compared with standard rehabilitation. *Am J Sports Med* 46(10):2432–2440
39. Lee DH, Kim JM, Lee BS, Kim KA, Bin SI (2012) Greater axial trough obliquity increases the risk of graft extrusion in lateral meniscus allograft transplantation. *Am J Sports Med* 40(7):1597–1605
40. Jeon B, Kim JM, Kim JM, Lee CR, Kim KA, Bin SI (2015) An osteophyte in the tibial plateau is a risk factor for allograft extrusion after meniscus allograft transplantation. *Am J Sports Med* 43(5):1215–1221
41. Jung Y, Choi N, Victoroff BN (2011) Arthroscopic stabilization of the lateral capsule of the knee in meniscal transplantation. *Knee Surg Sports Traumatol Arthrosc* 19(2):189–191
42. LaPrade CM, Ellman MB, Rasmussen MT, James EW, Wijdicks CA, Engebretsen L, LaPrade RF (2014) Anatomy of the anterior root attachments of the medial and lateral menisci: a quantitative analysis. *Am J Sports Med* 42(10):2386–2392
43. Bezerra FS, Alves JN, Silva MAS, Trajano ETL, Ferreira TA, Vasconcellos HA, Valença SS (2007) Quantitative and descriptive analysis of the meniscotibial ligament in human corpses. *Braz J Morphol Sci* 24(4):211–213

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.